										Required Prior to Kindergarten	Required Prior to 7th Grade	
		Vaccine	Birth	2 Months	4 Months	6 Months	12-15 Months	15-18 Months		4-6 Years	11-12 Years	13-18 Years
Required for Child Care		Haemophilus influenzae type b (HIB)		Hib	Hib	Hib	Hib					
		Pneumococcal (PCV)		PCV	PCV	PCV	PCV					
	Required for School	Hepatitis B (HepB)	НерВ	HepB		HepB						
		Diphtheria, Tetanus, Pertussis (DTaP)		DTap	DTaP	DTaP			age 2	DTaP		
		Poliovirus (Polio) (IPV)		IPV	IPV	IPV				IPV		
		Measles, Mumps, Rubella (MMR)					MMR		e by	MMR		
		Varicella (Chicken Pox) ¹					Varicella		dat	Varicella		
		Tetanus, Diphtheria, Pertussis (Tdap)							up-to		Tdap	
		Meningococcal ACWY (MCV4) ²							Children should be up-to-date by age		MCV4	MCV4 second dose, at age 16
		Meningococcal B (MenB) ³										MenB 2 doses, age 16-18
		Hepatitis A (HepA)					HepA	HepA				
	Recommended	Rotavirus (RV1 or RV5)		RV1/RV5	RV1/RV5	RV5						
		Human Papillomavirus (HPV)									HPV 2 or 3 doses ⁴	
		Influenza				Influenza Every flu season						
		COVID-19				COVID-19 Every respiratory virus season						
		RSV	RSV antibody for bables				Children 8-19 months with risk factors ⁵					

¹ Vaccine or documentation of history of disease.

VERMONT DEPARTMENT OF HEALTH

² Recommended for all. Required only for residential students entering 7th grade newly enrolled in 8 -12.

³ Recommendation for MenB vaccine is based on clinical discretion. Beginning at age 16, two doses at least one month apart.

⁴ If you start the series before age 15, only 2 doses are recommended. If you start after age 15 or are immunocompromised, then 3 doses are recommended.

⁵ Recommendation for children 8-19 months old who are at high risk for severe RSV during their second RSV season.